



# Spectrophotometric determination of Chlorpromazine Hydrochloride by oxidative coupling reaction with P-Bromo aniline using Ammonium Ceric Sulphate Dihydrate

Israa T. Humeidy

Department of Chemical Eng. / College of Engineering / University of Tikrit

[dr.israatalib@yahoo.com](mailto:dr.israatalib@yahoo.com)

Received date: 19 / 5 / 2015

Accepted date: 14 / 9 / 2015

## ABSTRACT

*A new sensitive spectrophotometric method for the determination of chlorpromazine hydrochloride (CPZH) was developed. The method is based on an oxidative coupling reaction with p-bromoaniline using ammonium ceric sulphate dihydrate as oxidizing agent to produce a violet color, soluble in water, stable product and absorbs at 553 nm. Beer's law was in the linear range 2.5 - 30 µg/ml of CPZH, the molar absorptivity, Sandell's sensitivity index and detection limit were  $6.5731 \times 10^3 \text{ liter. mol}^{-1}.\text{cm}^{-1}$ ,  $0.0541 \mu\text{g.cm}^{-2}$  and  $0.1361 \mu\text{g/ml}$  respectively. The RSD value was 1.20 - 1.36 % depending on the concentration. This method was applied successfully to the determination of chlorpromazine hydrochloride in two pharmaceutical formulations (tablets and injection).*

**Keywords:** spectrophotometric, CPZH, p-bromoaniline.

## التقدير الطيفي للكلوربرومازين بواسطة تفاعل الازدواج التأكسدي مع بارا-

### بروموانيلين بوجود كبريتات السيريوم الامونياكية

اسراء طالب حميدي

قسم الهندسة الكيماوية / كلية الهندسة / جامعة تكريت

[dr.israatalib@yahoo.com](mailto:dr.israatalib@yahoo.com)

تاريخ قبول البحث: ٢٠١٥ / ٩ / ١٤

تاريخ استلام البحث: ٢٠١٥ / ٥ / ١٩

#### الملخص



يتضمن البحث تطوير طريقة طيفية حساسة لتقدير عقار هيدروكلوريد الكلوربرومازين، الطريقة تستند على تفاعل الازدواج التأكسدي مع الكاشف بارا- بروموانيلين باستخدام العامل المؤكسد كبريتات السيريوم الامونياكية لتكوين ناتج بنفسجي اللون ذائب في الماء ويعطي أعلى امتصاص عند الطول الموجي ٥٥٣ نانوميتر. كانت حدود قانون بير في مدى التراكيز ٢.٥ - ٣٠ مايكروغرام/مل من هيدروكلوريد الكلوربرومازين. والامتصاصية المولارية  $6.5731 \times 10^3$  لتر.مول<sup>-1</sup>. سم<sup>-1</sup> ودلالة ساندل ٠.٠٥٤١ مايكروغرام. سم<sup>-2</sup>. وتراوحت قيمة الانحراف القياسي النسبي 1.2٠ - 1.36 % ، وحد كشف ٠.١٣٦١ مايكروغرام/مل. تم تطبيق هذه الطريقة بنجاح لتقدير هيدروكلوريد الكلوربرومازين في المستحضرات الصيدلانية حبوب وحقن.

الكلمات الدالة: الطريقة الطيفية، هيدروكلوريد الكلوربرومازين، بارا-بروموانيلين.

## 1. Introduction

The scientific name for the chlorpromazine hydrochloride (CPZH) is[1]:  
3-(2-Chloro-10H-phenothiazin-10-yl)-N,N-dimethylpropan-1-amine hydrochloride.

The chemical structure is:



Chlorpromazine hydrochloride (CPZH), white or almost white, crystalline powder, very soluble in water, freely soluble in ethanol. It decomposes on exposure to air and light[1]. Chlorpromazine is a psychiatric medication that belongs to the class of drugs called phenothiazine antipsychotics, it works by helping to restore the balance of certain natural substances in the brain[2]. It is used to treat various problems such as severe depression or behavioural disturbances and it is used to treat schizophrenia[3] and it also used to treat nausea, vomiting, severe pain, relieve prolonged hiccups, relieve restlessness/anxiety before surgery, and help treat tetanus[4]. side effects of chlorpromazine, HCl is low heat and drowsiness and nasal congestion and difficulty in urination and other side effects of low blood pressure and accelerated heart rate (in the case of intramuscular injection)[5].

Different analytical methods have been used for the determination of CPZH such as spectrophotometric methods[6-17], HPLC[18-24], gas chromatography[25-27], flow-injection techniques[28-34] and electrochemical method[35-36].

### Aim of research

In this research a rapid, sensitive and simple spectrophotometric method for determining of chlorpromazine hydrochloride (CPZH) in pure form as well as in pharmaceutical formulations based on the oxidative coupling using p-bromoaniline in presence of ammonium ceric sulphate dihydrate.

## 2. Experimental

### Apparatus

The apparatuses used shown in the Table (1).

**Table (1):** Apparatuses used by origin.

No.	Apparatus used by origin
1	Shimadzu UV/VIS 160 spectrophotometer, Japan
2	pH meter UK Jenway 3310
٣	Sartorius BL210 S AG Germany
٤	Ultrasonic with water bath, power sonic 410 Lab tech Korea
5	Hot Plate with Magnetic Stirrer (BIOSAN MSH 300)

### Reagents and chemicals used

All chemicals and analytical reagents used in this research are high purity and shown in the Table (2).

**Table (2):** Reagents and materials chemistry used.

Chemicals	Chemical structure	Company
Chlorpromazine, HCl	$C_{17}H_{19}ClN_2SHCl$	SDI
Ammonium ceric sulphate dihydrate	$Ce(NH_4)_4(SO_4)_4.2H_2O$	Fluka
p-Bromoaniline	$H_2SO_4$	BDH
Sulphuric acid	$C_6H_6BrN$	BDH
Sodium hydroxide	NaOH	BDH

### Preparation of solutions<sup>(37)</sup>

#### 1-Standard CPZH solution,250 µg/ml

This solution is prepared by dissolving 0.025 g of CPZH in amount of distilled water and the volume is diluted to 100 ml with distilled water in a volumetric flask. This solution is kept in a brown bottle, where it is stable for one week, at least.

#### 2- P- bromoaniline reagent solution ( $1 \times 10^{-2}M$ )

This solution is prepared by dissolving 0.1720m g of p-bromoaniline in 5 ml of ethanol and the volume is completed to 100 ml in a volumetric flask with distilled water.

### ٣- Ammonium ceric sulphate dihydrate solution ( $1 \times 10^{-2} M$ )

A 0.6320 g of ammonium ceric sulphate dehydrate ( $Ce^{+4}$ ) is dissolved in amount of distilled water and the volume is completed to 100 ml in a volumetric flask with distilled water.

### ٤-Sodium hydroxide solution (approximate,0.1 M )

This solution is prepared by dissolving 4.000 g of pure sodium hydroxide in 100 ml of distilled water.

### 5- Sulphuric acid solution (approximate,0.1M)

This solution is prepared by appropriate dilution of 0.5 ml of the concentrated sulphuric acid (18.289 M) solution to 100 ml with distilled water in a volumetric flask.

### ٦- Interference solutions 1000 $\mu g/ml$

A 0.1000 g of each foreign compounds is dissolved in distilled water then the volume is completed to 100 ml in a volumetric flask with distilled water.

### 7-Solution of CPZH tablets formulation (250 $\mu g/ml$ )

Largactil Chlorpromazine -HCl (100mg), Oubari Pharma - Aleppo-Syrie, every tablet contains 100 mg of CPZH and the solution is prepared as follows:

Ten tablets are weighed (4.1649g) and powdered well, then a weight of 0.1041 g of this powder is dissolved in an amount of distilled water, and then the solution filtered by paper filtration (604, RUNFILTER, Q240 mm), the volume is completed with distilled water in a volumetric flask of 100 ml , which gives the concentration of 250  $\mu g/ml$ .

### 8- Solution of CPZH injection formulation (25 $\cdot \mu g/ml$ )

Largactil Chlorpromazine -HCl (25mg/5ml), Oubari Pharma - Aleppo - Syrie, each injection contain of 25 mg of chlorpromazine and this solution is prepared by transferred of 50 ml of this formulation to volumetric flask and the volume was completed to 100 ml with distilled water to obtain a solution with a concentration of 250  $\mu g/ml$ .

### Preliminary Investigations

A 1.0 ml of p-bromoaniline reagent is added to 2.0 ml of standard CPZH solution in the presence of 0.5 ml of ammonium ceric sulphate dehydrate( $Ce^{+4}$ ) solution and diluted with distilled water in a 25 ml volumetric flask, a violet color product. Absorption spectrum of the colored dye against its corresponding blank reagents shows maximum absorption at 553 nm in contrast to blank reagent which shows no absorbance at this wavelength.

### 3. Results and Discussion

#### Optimization of the experimental conditions

To establish the optimum conditions, the effect of various variables on the intensity of the absorption was studied by adding 2.0 ml of standard CPZH solution (250 $\mu$ g/ml), 1.0 ml of p-bromoaniline and 0.5ml of  $Ce^{+4}$  and measuring the absorption at 553 nm versus the blank reagent.

#### Effect of acid

The effect of acid was studied by adding 0.5-3.5 ml of strong acids and weak acids (0.1 M). the results showed the  $H_2SO_4$  solution gives a higher absorption for colored product at a wavelength of 553 nm compared with other acids used , so this acid was chosen in subsequent experiments. the results are shown in Table (3).

**Table (3):** Effect of acid

Acid (0.1M)	ml of Acid /Abs.							
	With out	0.5	1	1.5	2	2.5	3	3.5
$H_2SO_4$	0.275	0.397	0.425	0.431	0.445	0.428	0.384	0.361
$HNO_3$	0.275	0.223	0.236	0.254	0.293	0.310	0.284	0.242
HCl	0.275	0.166	0.183	0.219	0.241	0.246	0.231	0.124
$CH_3COOH$	0.275	0.083	0.096	0.105	0.117	0.112	0.073	0.051

The results shown in the Table (3) indicate that the volume of 2.0 ml (0.1M) of sulfuric acid (pH= 1.8) is the optimum amount because of highest absorbance, so it was used in subsequent experiments.

### Effect of the amount of oxidizing agent

This study was conducted to select the best amount of oxidizing agent ( $Ce^{+4}$ ) ( $1 \times 10^{-2}$  M) by adding different volumes (0.3-2.5 ml) of oxidizing agent to volumetric flasks containing 2.0 ml of CPZH (250  $\mu$ g/ml) and 1.0 ml of the reagent solution ( $1 \times 10^{-2}$  M), then addition of 2.0 ml of 0.1 M sulfuric acid and the volume was completed to 25ml with distilled water, the results are shown in Table (4).

**Table (4):** Effect of the amount of oxidizing agent.

ml of $Ce^{+4}$ ( $1 \times 10^{-2}$ M)	0.3	0.5	0.7	1.0	1.2	1.5	1.7	2.0	2.5
Absorbance	0.337	0.445	0.458	0.473	0.441	0.326	0.273	0.246	0.216

The results shown in the Table (4) indicate that the volume of 1.0 ml of ( $Ce^{+4}$ ) ( $1 \times 10^{-2}$  M) is the optimum amount because of highest absorbance, so it was used in subsequent experiments.

### Effect of the amount of coupling reagent

The effect of the amount of coupling reagent was studied by adding different volumes (0.3-2.5ml) of p- bromoaniline reagent ( $1 \times 10^{-2}$  M) to the volumetric flasks containing 2.0 ml of CPZH (250 $\mu$ g/ml) and 1.0 ml of the ( $Ce^{+4}$ ) ( $1 \times 10^{-2}$  M), then the addition of 2.0 ml of 0.1 M sulfuric acid and the volume is completed to 25ml with distilled water, the results are shown in Table (5).

**Table (5):** Effect of the amount of coupling reagent.

ml of p-bromoaniline ( $1 \times 10^{-2}$ M)	0.3	0.5	0.7	1.0	1.2	1.5	1.7	2.0	2.5
Absorbance	0.359	0.416	0.451	0.474	0.485	0.481	0.473	0.452	0.436

It is clear that the volume of 1.2 ml of coupling reagent ( $1 \times 10^{-2}$  M) is the optimum amount because it gave the highest absorption. So it is adopted in subsequent experiments

### Effect of oxidation time

To a series of volumetric flasks, each containing 2.0 ml of CPZH (250  $\mu$ g/ml), 1.0 ml of ( $Ce^{+4}$ ) ( $1 \times 10^{-2}$  M) and the solutions were left for different periods of time, then 1.2 ml of p-bromoaniline reagent ( $1 \times 10^{-2}$  M) and 2.0 ml of 0.1 M sulfuric acid solution were added. The volume was completed to 25ml with distilled water, and the absorption of solutions was measured at a wavelength of 553 nm versus blank, the results are shown in Table (6).

**Table (6):** Effect of oxidation time.

Time(min)	5	10	15	20	25	30	35	40	45	50
Absorbance	0.429	0.467	0.486	0.485	0.485	0.483	0.482	0.481	0.479	0.479

Table (6) shows that 15 min is sufficient for the oxidation to be completed , so it is adopted in the subsequent experiments.

### Effect of temperature

The effect of temperature(5-60°C) on the absorption of the formed colored product were studied by using 2.0 ml of CPZH solution (250µg/ml) and 1.0 ml of (Ce<sup>+4</sup>) ( 1 × 10<sup>-2</sup>M), then 1.2 ml of p-bromoaniline reagent (1 × 10<sup>-2</sup>M) and 2.0 ml of 0.1 M sulfuric acid solution were added, then the volume is completed to 25 ml with distilled water in a volumetric flasks, and the absorption was measured at a wavelength of 553 nm versus blank reagent, the results are shown in Table (7). The optimum temperature is 25°C , so it is adopted in the subsequent experiments

**Table (7):** Effect of temperature.

Temp (°C)	5	10	15	20	25	30	35	40	45	50	60
Absorbance	0.294	0.329	0.386	0.409	0.485	0.483	0.467	0.385	0.311	0.269	0.175

### Effect of time on stability of the colored product

The stability time of the formed colored product was studied by taking 2.0 ml of CPZH (250 µg/ ml) with addition 1.0 ml of (Ce<sup>+4</sup>) (1×10<sup>-2</sup>M) and the solution is left for 15 min for oxidation, then 1.2 of p-bromoaniline (1 × 10<sup>-2</sup> M) and 2.0 ml of 0.1 M sulfuric acid solution were added,the volume is completed to 25 ml in a volumetric flasks with distilled water. It is observed that the absorption becomes constant directly after dilution and remain unaltered for 60 min. The results are shown on Table (8).

**Table (8):** Effect of time on stability of the colored product.

Time (min.)	Direct mixing time	5	10	15	20	25	30	35	40	45	50
Absorbance	0.485	0.484	0.485	0.385	0.485	0.486	0.485	0.484	0.484	0.483	0.482



### Effect of the solvents


The effect of the solvents on the formed colored product was studied, the dilution was carried out by different organic solvents instead of water. The results shown in Table (9) indicate that the water is a good medium for reaction and gives good absorption value at the wavelength of 553nm and due to its availability, it was used as the best solvent in the subsequent experiments.

**Table (9):** Effect of the solvents.

Solvent	Water	Acetone	DMSO	Methanol	Ethanol
Absorbance	0.485	0.426	0.257	0.195	0.216
$\lambda_{max}$ , nm	553	551	552	555	551

### Final absorption spectrum

The spectrum of the formed colored product by coupling of CPZH with p-bromoaniline ( $1 \times 10^{-2}M$ ) in the presence of  $Ce^{+4}$  ( $1 \times 10^{-2}M$ ) in acidic medium (pH 1.8) and temperature  $25^{\circ}C$  against its corresponding reagent blank show a maximum absorption at 553 nm in contrast to the blank reagent of zero absorbance at  $\lambda_{max}$ . The spectra are shown on Fig. (1).



**Fig.(1):** Final absorption spectrum of the determination CPZH.


**SB :** Absorption spectrum of CPZH solution versus blank reagent.

**SW:** Absorption spectrum of CPZH solution versus distilled water.

**BW:** Absorption of blank reagent versus distilled water.

### Procedure for construction of calibration curve

To a series of volumetric flasks (25ml), 0.25-3.0ml of (250 µg/ml) of CPZH were transferred, 1.0 ml of  $Ce^{+4}$  ( $1 \times 10^{-2}M$ ) was added and the solutions are left for 15 min for oxidation, then 1.2 ml of p-bromoaniline reagent ( $1 \times 10^{-2}M$ ) and 2.0 ml of 0.1M sulfuric acid solution (pH 1.8) were added at 25°C and the volumes are completed to the mark with distilled water. The absorbance is measured directly after dilution at 553 nm against the blank reagent. Fig. (2) illustrates that the calibration curve is linear over the concentration range of 2.5 - 30 µg/ml while higher concentrations show a negative deviation from Beer's law. The molar absorptivity value is  $6.5731 \times 10^3$  liter.  $mol^{-1}.cm^{-1}$  and the Sandell's sensitivity index  $0.0541 \mu g/cm^2$ .



**Fig.(2):** Calibration curve for determination CPZH by oxidative coupling with p-bromoaniline reagent.

### Accuracy and precision

Accuracy and precision were studied by measuring absorption (n=5) at 553 nm for two different concentrations of the drug within the limits of Beer's law, the average recovery (99.75 %) and the relative standard deviation (<1.39%) indicate that the method is of high accuracy and precision . The results are shown in Table (10).

**Table (10):** Results of accuracy and precision.

Conc. of CPZH(ppm)	RSD%	Recovery*%	Average recovery%	RE%
5	1.36	99.91	100.11	-0.04
10	1.20	100.30		0.30

\* Average of five determinations

### Detection limit

Detection limit was calculated by measuring the absorption for the lower concentration 2.5 µg/ml at optimal conditions (ten times) at 553 nm. The results are shown in Table (11).


**Table (11):** Detection limit.

Concentration µg/ ml	$\bar{X}$	S.D	D.L µg/ ml
2.5	0.158	0.0022	0.1361

\* Average of ten determinations


### The nature of the formed product

To know the nature of the formed violet color product (stoichiometry of drug with the reagent), Job's method and molar ratio method were applied. In both methods, the concentration of each of the standard CPZH solution and p-bromoaniline reagent solution is equal to  $7.0363 \times 10^{-4}$  M. In Job's method, in a series of volumetric flasks (25 ml ), different volumes of the drug solution ranging from 1-9 ml and different volumes (9-1 ml) of reagent solution were mixed. A 1.0 ml of  $(Ce^{+4})(1 \times 10^{-2}$  M) and 2.0 ml 0.1 M of sulfuric acid solution were added and volumes were completed to the mark with distilled water. The absorbance was measured at 553 nm against the blank reagent. The results Fig. (3) show that the ratio is 1:1.



**Fig. (3):** Job's method of formed product by oxidative coupling of CPZH with p-bromoaniline reagent.

In molar ratio method, 1 ml of the standard drug solution in a series of volumetric flasks (25 ml) were transferred and different volumes 0.2-3.5 ml of N,N-DMPPDADH reagent solution, 1.0 ml of  $(Ce^{+4})$  ( $1 \times 10^{-2}M$ ) and 2.0 ml 0.1 M of sulfuric acid solution were added. The volumes were completed to the mark with distilled water and the absorbance was measured at 553 nm against the blank reagent. Molar ratio was found to be 1:1. The results are shown in Fig.(4) which is in agreement with the Job's method results.



**Fig. (4):** Molar ratio method for the product formed by oxidative coupling of CPZH with p-bromoaniline reagent.


### Effect of interferences

In order to test the efficiency and selectivity of the proposed method, the effect of some foreign substances (lactose, fructose , maltose, starch and glucose) that usually present in dosage forms are studied by taking volumetric flasks ( 25 ml) containing 2.0 ml of CPZH (250  $\mu g$  /ml), then different volumes ( 2.5, 5.0 ,7.5 ml) of foreign substances (1000  $\mu g$ /ml) have been added resulting in a final concentration of (100, 200 ,300  $\mu g$ /ml ). The optimum conditions have been applied and the volumes have been completed to the mark with distilled water. The absorbance was measured at 553 nm versus blank reagent and recovery is calculated. The results showed that there is no interferences Table (12).

**Table (12):** Effect of interferences.

Foreign compound	Recovery (%) of 20 µg/ml of CPZH per µg /ml foreign compound added		
	100	200	300
<b>lactose</b>	<b>100.08</b>	<b>100.27</b>	<b>99.76</b>
<b>Fructose</b>	<b>98.48</b>	<b>99.35</b>	<b>97.55</b>
<b>Maltose</b>	<b>97.39</b>	<b>98.66</b>	<b>98.90</b>
<b>Starch</b>	<b>99.89</b>	<b>97.51</b>	<b>99.43</b>
<b>Glucose</b>	<b>100.47</b>	<b>99.52</b>	<b>100.12</b>

The proposed equation for reaction can be written as follows:



### Applications

This method was applied for the determination of CPZH in its pharmaceutical formulations (tablets and injection).

### Direct method

In this method, different volumes (0.25, 0.5ml) of a pharmaceutical formulation solutions (250 µg/ml) were transferred to 25 ml volumetric flasks and the resulting concentrations (2.5, 5 µg/ml) and were treated as in construction of calibration curve. The absorbance was measured at 553 nm for five times. Recovery and RSD were calculated and the results are shown in Table (13).

**Table (13):** Determination of CPZH in pharmaceutical formulation


pharmaceutical formulation	CPZH present $\mu\text{g/ml}$	CPZH measured $\mu\text{g/ml}$	RSD,%	Recovery *,%
Injection largapromatctil chlorpromazine -HCl 25 mg/5ml	2.5	2.49	1.44	99.60
	5.0	5.02	1.24	100.40
Tablets largapromatctil chlorpromazine -HCl 100 mg	2.5	2.52	1.56	100.8
	5.0	4.96	1.53	99.20

\* Average of five determinations


Table (13) shows the efficiency and success of the developed method for the determination of CPZH in its pharmaceutical formulation.

#### standard additions method

To prove that the developed method is free from interferences , method of standard additions is applied for determining of CPZH in its pharmaceuticals. Different volumes (0.25,0.5 ml) of a pharmaceutical formulation solutions (250  $\mu\text{g/ml}$ ) were transferred to six volumetric flasks (25 ml) for each volume, then increasing volumes (0.25-2.0 ml) of 250  $\mu\text{g/ml}$  of CPZH standard solution were added with leaving the sixth flask without addition. The solution was treated as in construction of calibration curve. The absorbance were measured at 553 nm Fig.(5,6) the measured concentration was calculated from the equation of the straight line and the results of Recovery and RE shown in the Table (14).



**Fig.(5)** Standard additions curve for the determination of CPZH in injection



**Fig.(6)** Standard additions curve for the determination of CPZH in tablets.

**Table (14):** Results of standard additions method.

pharmaceutical formulation	CPZH present $\mu\text{g/ml}$	CPZH measured $\mu\text{g/ml}$	Recovery <sup>*</sup> , %
Injection largapromatcil chlorpromazine -HCl 25mg /5ml	2.5	2.51	100.40
	5.0	5.01	100.20
Tablets largapromatcil chlorpromazine -HCl 100mg	2.5	2.49	99.60
	5.0	4.99	99.80

The results shown in table 12 indicate that method of standard additions is in agreement with the direct method within the acceptable range of error, indicating that the method is satisfactory and free from interferences.

#### 4. Conclusions

The results obtained confirm that the proposed method is simple, rapid and of good sensitivity for the determination of chlorpromazine hydrochloride CPZH. The method is based on oxidative coupling between CPZH and p-bromoaniline reagent in presence of ammonium ceric sulphate dehydrate ( $\text{Ce}^{+4}$ ) in acidic medium to form a violet colored dye which is water soluble, stable and shows a maximum absorption at 553 nm. This method does not require temperature control, nor use of organic solvents, or solvent extraction and it can be applied successfully for determination of CPZH in pharmaceuticals formulation.

#### References

- [1] British Pharmacopeia, 6<sup>th</sup> Ed., by system simulation Ltd., the stationary office, London, (2009). in CD-ROM"
- [2] K. Heinrich, ;*Psychopharmaka in Klinik and Praxis*; 2<sup>nd</sup> edn., Germany, 1983, P 432.
- [3] J. Ross and I. Tarazi ;*Pharmacotherapy of Psychosis and Mania*; 11<sup>th</sup> Ed., New York, 978-981, (2006).
- [4] J. Applphysiol. ;*Appl.phys. Lett.*; 50(3), 509-512 , (1981).
- [5] Martindale, ;*The extra Pharmacopoeia*; Pharmaceutical Society of Great Britain, 28<sup>th</sup> Ed., 1509-1615 , (1982) .



- [6] K. Basavaiah; *Indirect spectrophotometric determination of some biologically important phenothiazines using potassium dichromate, iron(II) and 1,10-phenanthroline*; Indian Journal of Chemical Technology, 11, 632-638, (2004).
- [7] K. Basavaiah, P. Nagegowda, H. C. Pramwula and B. C. Somashekar; *Sensitive titrimetric and spectrophotometric assay methods for chlorpromazine with bromate – bromide mixture and two dyes*; Indian Journal of Chemical Technology, 12, 25-29, (2005).
- [8] S. M. Al-Talib and T. N. Al – Sabha; *Spectrophotometric determination of some phenothiazines using N-chlorosuccinimide*; Jou. Raf. Sci., 20(4), 27-37, (2009).
- [9] A.S. AL-Ayash , F. Jasim and W. Alwan; *Spectrophotometric determination of chlorpromazine hydrochloride in pharmaceutical preparations*; National Journal of Chemistry, 33, 42-53, (2009).
- [١٠] W. A. Abdullah; *Spectrophotometric determination of chlorpromazine hydrochloride using oxidation reduction method based on reaction of iron (II) with 2, 2'-bipyridyl reagent*; Jou. Raf. Sci., 22(2), 79-93, (2011).
- [١١] G.V. Raju, S.K. Jaffar, M. P. priya , S.M. Ibrahim, P. Rajakumar , E.Muralinath and M.Guru Prasad; *Application of MBTH/FeCl<sub>3</sub>: A reliable and coast effective analytical methodology for routine pharmaceutical determination of chlorpromazine hydrochloride*; IJRRPAS, 1(4), 202-206, (2011).
- [١٢] Z. Yi-mi, L. Ke-ke, Y. Cun-ling and W. Zhi-ke; *Spectrophotometric determination of chlorpromazine hydrochloride by 1,10-phenanthroline*; Chinese Journal of Analysis Laboratory, 01, 39-41, (2012).
- [١٣] M.E.M. Hassouna , A.M. Adawi, E.A. Ali; *Extractive spectrophotometric determination of chlorpromazine and trifluoperazine hydrochloride in pharmaceutical preparations*; Egyptian Journal of Forensic Sciences, 2 (2), 62–68, (2012).
- [١٤] M.J.H.AL-Kaffiji and A..M.S..AL-Anbakey; *New chromogenic reagent for the spectrophotometric determination of chlorpromazine HCl in aqueous solutions and pharmaceutical formulations*; Int. J. Pharm. Pharm. Sci., 5(3), 606-611, (2013).
- [١٥] N. K. A. Al-Hayani, and F. K. Mohammad; *A simple spectrophotometric assay for stability determination of chlorpromazine in veterinary injectable solutions*; International Journal of the Bioflux Society, 5(2), 66-69, (2013).





- [١٦] M. M. Al-Rufaie, A. N. Al-Sharefy and K. H. kadim; *Spectrophotometric determination of chlorpromazine hydrochloride in pharmaceutical preparations by using oxidativ coupling reaction*; Iraqi National Journal of Chemistry, 51, 338-347, (2013).
- [1٧] L. A. Al Shatti; *Method development and validation of assay of chlorpromazine hydrochloride tablet formulation using ultra violet visible spectrophotometry*; J. Anal. Bioanal. Tech., 5(2), 2-4, (2014).
- [1٨] T. Ohkubo, R. Shimoyama, K. Sugawara; *Determination of chlorpromazine in human breast milk and serum by high-performance liquid chromatography*; J. Chromatogr., 614(2), 328-332, (1993).
- [1٩] M. D. Rukhadze, V. M. Okudzhava, M. M. Aleksishvili, S. K. Tsagareli and T. G. Makharadze; *Determination of chlorpromazine in blood serum by ion-pair reversed-phase HPLC: Chlorpromazine pharmacokinetics in rabbits*; Pharmaceutical Chemistry Journal, 33(9), 502-504, (1999).
- [٢٠] C. Pistos and J. T. Stewart; *Direct injection HPLC method for the determination of selected phenothiazines in plasma using a Hisep column*; Biomedical Chromatogr., 17(7), 465-470, (2003).
- [2١] P. Shetti and A. Venkatachalam; *Stability indicating HPLC method for simultaneous quantification of trihexyphenidyl hydrochloride, trifluoperazine hydrochloride and chlorpromazine hydrochloride from tablet formulation*; E-Journal of Chemistry, 7(1), 299-313, (2010).
- [2٢] S.Venkatesh, M.B.Kumar, S.Ramachandran and G.N.Sameer; *HPLC method development, valdation and its application to stability studies of chlorpromazine hydrochloride tablets*; IRJP, 1(1), 225-232, (2010).
- [2٣] J.M. Dhabab; *Simultaneous determination of chlorpromazine and trifluoperazine in pharmaceutical preparations using high performance liquid chromatography*; Al-Mustansiriya J. Sci, 22(2), 123-128, (2011).
- [2٤] N.U. Rani, K. Divya, G. Sahithi; *New validated RP-HPLC method for simultaneous estimation of chlorpromazine and trihexyphenidyl HCl in tablets*; IJAPA, 4(4), 134-137, (2014).
- [2٥] L.D. Gruenke, J.C. Craig, F.D. Klein, T.L. Nguyen, B.A. Hitzemann, Holaday, H.H. Loh, L. Braff, A. Fischer and I.D. Glick; *Determination of chlorpromazine and its major*



- metabolites by gas chromatography/mass spectrometry: application to biological fluids*; Biomed. Mass Spectrom., 12(12),707-713, (1985).
- [2٦] R. Ninci, M.G. Giovannini, L. D. Corte and G. Sgaragli; *Isothermal gas chromatographic determination of nanogram amounts of chlorimipramine, chlorpromazine and their N- desmethyl metabolites in plasma using nitrogen- selective detection*; J. Chromatogr., 381(2),315-322, (1986).
- [2٧] L. Zhang, P. Wu, Y. Zhang, Q. Jin, D. Yang, L. Wang, and J. Zhang; *A GC/MS method for the simultaneous determination and quantification of chlorpromazine and diazepam in pork samples*; Anal. Methods, 6(2), 503-508, (2014).
- [2٨] D. Chen, A. Rios, M.D. Castro and M. Valcarcel; *Simultaneous flow- injection determination of chlorpromazine and promethazine by photochemical reaction*; Talanta, 38(11),1227-1233, (1991).
- [2٩] T. Ghous and A. Townshend; *Flow injection determination of chlorpromazine by inhibition of glutamate dehydrogenase*; Analytica. Chimica. Acta., 387(1), 47–51, (1999).
- [٣٠] D. Daniel and I.R.G. Gutz; *Spectroelectrochemical determination of chlorpromazine hydrochloride by flow-injection analysis*; J. Pharm. Biomed. Anal., 37(2),281-286,(2005).
- [31] M. G. F. Sales , J. F.C. Tomas and S. R. Lavandeira; *Flow injection potentiometric determination of chlorpromazine*; Journal of Pharmaceutical and Biomedical Analysis, 41(4), 1280–1286, (2006).
- [32] J. A. Ortuno, A.Gil and C. Sanchez-Pedreno; *Flow-Injection coulometric detection based on ion transfer and its application to the determination of chlorpromazine*; Sensors, 8, 3678- 3688, (2008).
- [33] I.M.A.Shakir and M.J.H. Al-kaffiji; *Flow-injection spectrophotometric determination of chlorpromazine HCl based on releasing of sodium persulphate from hydrogel beads, study and applications*; Wasit Journal for Science & Medicine, 7(3), 110-125, (2014).
- [34] I.M.A.Shakir and M.k.Hammood; *New turbidimetric-continuous flow injection analysis method for the determination of chlorpromazine HCl in pharmaceutical preparation using linear array ayah 5SX1-T-1D-CFI analyser*, Iraqi Journal of Science, 55(2),594-605, (2014).
- [35] N. Yongnian, L. Wang and S. Kokot; *Voltammetric determination of chlorpromazine hydrochloride and promethazine hydrochloride with the use of multivariate calibration*; Analytica. Chimica. Acta. , 439(1), 159-168, (2001).

[36] E. Y.Z. Frag, M.A. Zayed, M.M. Omar, S. E.A. Elashery and G.G. Mohamed; *Potentiometric determination of chlorpromazine HCl using carbon paste electrode in pure and pharmaceutical preparations*; Int. J. Electrochem. Sci., 7, 650 – 662, (2012).

[37] D.A. Skoog , D.M. West,F.J.Holler ,and S.R.Crouch ; *Fundamentals of Analytical Chemistry*; 8th Ed. Brooks/Cole- Thomson Learning, Belmont, 71-86,(2004).

#### AUTHOR



**Israa Talib Humeidy** : Instructor in Analytical Chemistry Department of Chemical Engineering/College of Engineering / University of Tikrit  
B. Sc. in chemistry from College of Sciences/ Mosel University/Iraq  
M. Sc. In Analytical Chemistry from College of Education Pure Sciences/  
Tikrit University/ Iraq Ph. D In Analytical Chemistry from College of  
Sciences / Tikrit University/ Iraq.