



**SYNTHESIS, CHARACTERIZATION OF SOME METAL
COMPLEXES OF DIETHANOLAMINEDITHIOCARBAMATE AND
(1,10-PHENATHROLINE)**

Hasan .A. Mohammed*

Department of Chemistry, College of science, university of Kirkuk

**SYNTHESIS, CHARACTERIZATION OF SOME METAL COMPLEXES OF
DIETHANOLAMINEDITHIOCARBAMATE AND (1,10-PHENATHROLINE)**

Hasan .A. Mohammed*

Department of Chemistry, College of science, university of Kirkuk

* Hassanahmed88@yahoo.com

* Hassanahmed88@yahoo.com

ABSTRACT

A dithiocarbamate ligand; sodiumdiethanolaminedithiocarbamate Na(deadtC) is synthesized from the reaction of diethanolamine with carbon disulfide and sodium

hydroxide. (phen)= 1,10-phenanthroline . Addition of metal salts, gave complexes of the types: $[M(\text{deadtC})_2 \text{ phen}]$, where $M = \text{Fe(II), Co(II), Ni(II) and Cu(II)}$. $[M(\text{deadtC})_3]$ where $M = \text{Fe(III)}$. The ligand (deadtC) behaves as a bidentate and coordinated to the metal ion centers either through the sulfur atom of its dithiocarbamate and or through the nitrogen atoms of the phenanthroline. All the synthesized ligands and complexes are characterized by elemental analyses,



conductivity, infrared, electronic spectra and susceptibility measurements. From the obtained data octahedral geometry for the complexes have been suggested.

Keywords: diethanolamine , dithiocarbamate, 1,10-phenanthroline, Transition metal complexes.

تحضير و تشخيص بعض معقدات ثنائي ايثانول امين ثنائي الكبريت كاربميت مع (١٠, ١) فينانثرولين .

*حسن أحمد محمد

جامعة كركوك – كلية العلوم – قسم الكيمياء

* Hassanahmed88@yahoo.com

الملخص:

ان الليكاند ثنائي الكبريت كاربميت : ثنائي ايثانول امين ثنائي الكبريت كاربميت (*deadtc*) والمحضر من ثنائي ايثانول امين مع كبريتيد الكربون وهيدروكسيد الصوديوم. اضافة املاح الفلزات تعطي معقدات ذات الصيغ $[M(\text{deadtc})_2 \text{phen}]$ حيث $M = \text{Fe(II)}, \text{Co(II)}, \text{Ni(II)}, \text{Cu(II)}$ ، $(\text{phen}) = 1, 10$ -فينانثرولين ، $[M(\text{deadtc})_3]$ حيث $M = \text{Fe(III)}$ تم تشخيص الليكاند والمعقدات المحضرة بواسطة التحليل



الدقيق للعناصر ، التوصيلية ، الاشعة تحت الحمراء ، الاطياف الالكترونية والحساسية المغناطيسية بينت القياسات ان الليكاند يناسق بشكل ثنائي السن من خلال ذرتي الكبريت لثنائي كبريت الكربميت ومن خلال ذرات النتروجين للفينوثرولين وان الشكل الهندسي المقترح للمعقدات هو ثماني السطوح

الكلمات المفتاحية: ثنائي ايثانول امين، ثنائي الكبريت كاربميت ، ١, ١٠- فيناثرولين ، املاح الفلزات.

1. INTRODUCTION

Metal complexes of dithiocarbamates have been widely studied in recent years [1–5], because of their wide range of applications in agriculture as pesticides, in medicines, in industry as vulcanization accelerators, Besides that, nitrogen donor adducts of dithiocarbamate complexes are also widely used in the preparation of thin semiconductor [6–8]. Recently, 1,1-dithiolate ligands have attracted much attention, mainly because of interesting photophysical properties derived from their extensive electron delocalization over all of the ligand atoms [9]. Understanding bonding in these complexes will allow the design of new effective complexes used in biological systems [10]. Dithiocarbamate ligands display a strong propensity of binding to metal atoms such as nickel [11].

2. Experimental

2.1 Materials and Methods

Higher grade reagents and solvents were commercially available (Fluka A.G., Merck, BDH) and used as received. Infrared spectra were recorded on a Nicolet 100 FTIR



spectrophotometer in the 400-4000 cm^{-1} range using KBr discs. NMR spectroscopy recorded by Av 300 instrument. Conductivity measurements were carried out on 10^{-3} M solution of the complexes in DMF using conductivity meter Jenway PCM3 at an ambient temperature. The electronic spectra were recorded on a PgT92+ UV-visible spectrophotometer for 10^{-3} M solutions of complexes in DMF as solvent at 25°C using 1 cm quartz cell. Melting points were recorded on an Electrothermal 9300 apparatus. The magnetic susceptibility measurements were carried out at 25°C on the solids by Gouy's method using Sherwood Scientific instrument.

2.2 Syntheses of the ligand (sodium diethanolamine dithiocarbamate)

Na(deadtC)

Diethanolamine (1.05g, 0.01 mol) was dissolved in 30 ml of water containing sodium hydroxide (0.4 g, 0.01 mol) with constant stirring. The resulting solution was cooled in an ice bath, and carbon disulfide (0.76 g, 0.01 mol) was added dropwise with stirring. The mixture was stirred for 30 min. The yellow precipitate formed, was filtered off, washed with diethyl ether and dried in vacuum.

2.3 Synthesis of the complexes [M(deadtC)₂ phen] M= Fe(II), Co(II), Ni(II),

Cu(II)

A solution of sodium diethanolaminedithiocarbamate (2.03 g, 0.01 mol) in 25 ml of water was added dropwise to each aqueous solution of $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ (0.99 g, 0.005 mol) and $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (1.15g, 0.005 mol) and $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (1.18 g, 0.005 mol) and $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ (0.845 g, 0.005 mol) with a constant stirring at a room temperature followed by addition of (0.005 mol) of (1,10 - phenathroline). After 30 min, the precipitate formed was filtered off, washed with 10ml of water and dried under vacuum.


2.4 Synthesis of the complexes [Fe(deadtC)₃]



A solution of sodium diethanolaminedithiocarbamate (2.03 g, 0.01 mol) in 25 ml of water was added to aqueous solution of $\text{FeCl}_3 \cdot 4\text{H}_2\text{O}$ (0.6984 g, 0.003 mol) dropwise with a constant stirring at a room temperature. After 30 min, the precipitate formed was filtered off, washed with 10 ml of water and dried under vacuum.

3 Results and Discussion

The ligand was prepared by the reaction of carbon disulfide with the secondary amine diethanolamine, in the presence of sodium hydroxide and the complexes were prepared by direct addition of the aqueous solution of the chloride salts of the studied metals to the ligand solution in aqueous media, using 1:2:1 (metal: ligand: adduct) and 1:3(metal: ligand) molar ratio. (Scheme 1).





Scheme (1) preparation of ligand and dithiocarbamate complexes

The composition and analytical data for all complexes are given in Table 1. The low values of molar conductivities of the complexes in DMF ($\approx 10^{-3} \text{ ohm}^{-1}.\text{cm}^2.\text{mol}^{-1}$) indicate that all the prepared complexes are non-electrolytes. [13]

Table 1: Analytical and some physical properties of the prepared complexes.

Comp.no.	Chemical formula	color	m.p °C	Λ $\text{ohm}^{-1}.\text{cm}^2.\text{mol}^{-1}$	Yield %	μ_{eff} B.M
1.	Na(deadc)	yellow	157	0.7	87	-----
2.	[Fe(deadc) ₂ phen]	red	192-193	12.0	70	4.90
3.	[Co(deadc) ₂ phen]	orange	181-182	14.7	82	4.78
4.	[Ni(deadc) ₂ phen]	Light green	215-216	9.2	90	2.90
5.	[Cu(deadc) ₂ phen]	Brown	245-246	11.8	84	1.92
6.	[Fe(deadc) ₃]	Brown	212-113	10.3	77	0.9



3.1 Electronic spectra

The band at 37313 cm^{-1} is assigned to $(\pi \rightarrow \pi^*)$ absorption for ligand [14]. The electronic spectra of the prepared iron (II) complex displays bands at (10365 cm^{-1}) refer to $({}^5T_{2g} \rightarrow {}^5E_g)$ and other charge transfer bands at $(30894-35460)$. The cobalt(II) complex displays bands at (10728 cm^{-1}) , (15635 cm^{-1}) and (27821 cm^{-1}) referring to ${}^3A_{2g}(F) \rightarrow {}^3T_{2g}(F)$, ${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(F)$, ${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(P)$ respectively and other charge transfer bands at $(30120-34965\text{ cm}^{-1})$. The nickel(II) complex displays bands at (11532 cm^{-1}) , (15013 cm^{-1}) and (23342 cm^{-1}) referring to $({}^3A_{2g}(F) \rightarrow {}^3T_{2g}(F))$, $({}^3A_{2g}(F) \rightarrow {}^3T_{1g}(F))$, $({}^3A_{2g}(F) \rightarrow {}^3T_{1g})$ respectively and other charge transfer bands at $(34873\text{ cm}^{-1}-35258\text{ cm}^{-1})$. The electronic spectrum of Cu(II) complex shows a band at (11247 cm^{-1}) , referring to $({}^2E_g \rightarrow {}^2T_{2g})$ and another charge transfer band at (28472 cm^{-1}) . [15] the electronic spectrum of Fe(III) shows a band at (31737 cm^{-1}) , which is assigned to charge transfer.

3.2 Magnetic susceptibility

The magnetic moments (μ_{eff}) for Fe (II), Co (II), Ni(II) and Cu(II) complexes are $(4.90, 4.78, 2.95, 1.95\text{ B.M})$ respectively suggest an octahedral geometry. [16- 17]. The higher magnetic moment values than spin only moment of Fe(II) and Co(II) complexes are due to orbital contribution. The magnetic moment of the Fe (III) complex (5.9 B.M) Suggests a high spin octahedral geometry [18].



Table 2: The electronic data for sodium diethanolamine dithiocarbamate Na(deadc) and its metal complexes.

No.	Compounds	Band absorption	Assignment
		cm ⁻¹	
1.	Na(deadc)	37313	$\pi \rightarrow \pi^*$
2.	[Fe(deadc) ₂ phen]	10365	$^5T_{2g} \rightarrow ^5E_g$
		35460	Charge transfer
3.	[Co(deadc) ₂ phen]	10728,15635,27821	$^3A_{2g}(F) \rightarrow ^3T_{2g}(F)$, $^3A_{2g}(F) \rightarrow ^3T_{1g}(F)$, $^3A_{2g}(F) \rightarrow ^3T_{1g}(P)$
		30120	Charge transfer
4.	[Ni(deadc) ₂ phen]	11532 ,15013,23342	$^3A_{2g}(F) \rightarrow ^3T_{2g}(F)$, $(^3A_{2g}(F) \rightarrow ^3T_{1g}(F))$, $(^3A_{2g}(F) \rightarrow ^3T_{1g})$
		34873	Charge transfer
5.	[Cu(deadc) ₂ phen]	11247	$^2E_g \rightarrow ^2T_{2g}$
		28472	Charge transfer
6.	[Fe(deadc) ₃]	31737	Charge transfer

3.3 Infra-Red Spectroscopy



The most important bands in the FT-IR spectra of the prepared ligand and the complexes are recorded in Table 2. For dithiocarbamate compounds, three important regions of IR spectra are of interest. These are associated with the stretching vibration of ν (N–C), ν (–C=S) and ν (M–S) and the single absorption band in the second region suggests a bidentate behavior [19,20]. The ligand show a strong absorption at (1480) cm^{-1} , which is assigned with the C-N stretching frequency, while the complexes showed absorption at (1484 -1509 cm^{-1}). They are assigned to ν (C-N) stretching frequency. Compared with the ligand, the complex to shift to a higher frequency by (8-46) cm^{-1} . The presence of a single strong band (955-1074) cm^{-1} due to a ν (CSS) mode in the spectra of the complex is strongly indicative of the bidentate behavior of the dithio ligand in the complexes. [21] the absorption bands at (410-486) cm^{-1} are assigned to ν (M-S) and ν (M-N).the absorption broad bands at (3383) cm^{-1} suggests (O-H) stretching frequency in the ligand and complexes. This band is unchanged or slightly shifted to higher frequency in the spectra of the complexes indicating that the OH group is not involved in coordination but engaged in intramolecular hydrogen bonding. [22]



Table (3): Selected IR bands of the sodium diethanolamine dithiocarbamate Na(deadc) and its metal complexes (cm⁻¹).

No.	Compounds	$\nu(\text{H-O})$	$\nu(\text{C=S})$	$\nu(\text{C-N})$	$\nu(\text{N-M})$	$\nu(\text{S-M})$
1.	Na(deadc)	3338	982	1480		
2.	[Fe(deadc) ₂ phen]	3386	1006	1490	568	468
3.	[Co(deadc) ₂ phen]	3342	995	1492	572	466
4.	[Ni(deadc) ₂ phen]	3347	991	1511	586	484
5.	[Cu(deadc) ₂ phen]	3280	993	1489	548	468
6.	[Fe(deadc) ₃]	3343	987	1508		495


3.4 NMR Spectroscopy

The NMR spectra for the ligand and [Cu(deadc)₂ phen] complex recorded in Av 300 spectrometer . The ¹H NMR spectra for the ligand and complex show signals at δ (4.08 , 3.54 , 2.7) and (9.02 , 8.50 , 8.00 , 7.7 , 4.9 , 3.7, 3.4) respectively. The signals at δ (8- 9.02) are assigned to (1,10-phenanthroline). The ¹³C NMR spectra show signals at δ (210 , 204) assigned to CS₂ in the ligand and complex as in Table (4).




Table (4): Important signals of ¹HNMR and ¹³C NMR Spectra of the Ligand and Complex.


No.	Compound	NMR	δ ppm
1.	Na(deadtC)	¹ HNMR	4.08(2H), 3.54(4H), 3.62(4H),2.7(4H)
		¹³ CNMR	39.2 ,55.4 , 55.7 ,59.7 , 210 (NCS ₂).
2.	[Cu(deadtC) ₂ phen]	¹ HNMR	3.4 (8H), 3.7 (8H), 4.9(4H),7.7(2H), 8.0(2H), 8.5(2H), 9.02(2H).
		¹³ CNMR	39.8 ,41.7 ,51.6 ,58.1, 105 , 138.9, 158.3 ,204.9(NCS ₂).




Scheme (2) The ¹³C NMR of sodium diethanolamine dithiocarbamate ligand



Scheme (2) The ¹HNMR of sodium diethanolamine dithiocarbamate ligand



Scheme (2) The ¹HNMR of [Cu(deadt)₂ phen] complex



Scheme (2) The ¹HNMR of [Cu(deadtC)₂ phen] complex

3 Conclusion

From the above discussion we suggested that the structure of (2,3,4,5) complexes are distorted octahedral except Fe (III) complex is octahedral. The mixed ligands effect to convert the symmetry of complex O_h to D_{4h}.



References

- [1] Lan-Feng Hou, Yun Zhong, Yan Meia and Jun Fanb “Bis[N,N-bis(2-hydroxyethyl)dithiocarbamato]copper(II)” Acta Cryst. E65, m1694,(2009).
- [2] P. Balaramesh, A. Abdul Jabbar and P.Venkatesh“Studies on stable platinum(II)metal complexes of ethyenediamine and diethanolamine dithiocarbamates” International Journal of Scientific and Research Publications, Volume 4, Issue 5, (2014)
- [3] G. Hogarth, Ebony-Jewel C.-R.C.R. Rainford-Brent, I. Richards “Functionalized dithiocarbamate complexes: Synthesis and molecular structuresof bis(2-methoxyethyl)dithiocarbamate complexes $[M\{S_2CN(CH_2CH_2OMe)_2\}_2]$ (M = Ni, Cu, Zn) and $[Cu\{S_2CN(CH_2CH_2OMe)_2\}_2][ClO_4]$ ”, Inorganica Chimica Acta ,362, 1361–1364,(2009)
- [4] H. Nabipour “Synthesis of a new dithiocarbamate cobalt complex and its nanoparticles with the study of their biological properties” Int.J.Nano.Dim 1(3): 225-232, (2011)
- [5] N.Srinivasan,P. Valarmathi,S. Thirumaran , S. Ciattini “Synthesis and spectral studies on NiS₄, NiS₂PN, NiS₂P₂ chromophores: Single-crystal X-ray structure of $[Ni(dbpdtc)_2]$ (dbpdtc = benzyl(4-(benzylamino)phenyl)dithiocarbamate)” Transition Met Chem vol35,815–819,(2010)
- [6] P.V. Subha a, P. Valarmathi a, N. Srinivasan a, S. Thirumaran a,* , K. Saminathan “Effect of size of metal ion on MS₄N₂ chromophore: Synthesis, spectral and single crystal X-ray structural studies on (2,20-bipyridine)bis(N-cyclohexyl-N-methyldithiocarbamato)M(II) (M = zinc, cadmium)” Polyhedron vol 29 ,1078–1082,(2010)



- [7] G. J. Perpetuo, M. R.L. Oliveira, J. Janczak, H. P. Vieira, F. F. Amaral, V. M. De Bellis “Syntheses, crystal structure and spectroscopic characterization of novel N-R-sulfonyldithiocarbamate zinc(II) complexes” *Polyhedron* vol 22 , 3355–3362,(2003)
- [8] D. C. Onwudiwe and P. A. Ajibade “Synthesis, Characterization and Thermal Studies of Zn(II), Cd(II) and Hg(II) Complexes of N-Methyl-N-Phenyldithiocarbamate: The Single Crystal Structure of $[(C_6H_5)(CH_3)NCS_2]_4Hg_2$ ” *Int. J. Mol. Sci.* 12, 1964-1978,(2011)
- [9] G. Marimuthu, K. Ramalingam, C. Rizzoli “Synthesis, spectral, thermal and BVS investigations on ZnS_4N^N/N coordination environment: Single crystal X-ray structures of bis(dibenzoyldithiocarbamato)(N^N)Zinc(II) complexes ($N^N = 1,10$ -phenanthroline, tetramethylethylenediamine and bipyridine)” *Polyhedron*, vol 29 1555–1560,(2010)
- [10] S. P. Sovilj , N. Avramovic, N. Katsaros “Syntheses and properties of mixed dinuclear copper(II) complexes with heterocyclic dithiocarbamates and a cyclic octadentate tertiary amine” *Transition Metal Chemistry* 29: 737–742, (2004).
- [11] J .Heard, *Prog Inorg Chem* 53:1,(2005)
- [12] M. Sarwar , S. Ahmad, S. Ahmad, S. Ali, S. Ahmed Awan “Copper(II) complexes of pyrrolidine dithiocarbamate” *Transition Metal Chemistry* vol 32:199–203(2007)
- [13] W.J. Geary, *Coord. Chem. Rev.* 7, 81(1971).
- [14] J.Cookson ,A.L.Emma , P. M.John , J. S.Christopher , L. P.Rowena , R. C.Andrew , G.B.Michael, and D.Paul, “Metal-directed assembly of large dinuclear copper (II) dithiocarbamateMacrocyclic complexes” *InorganicaChimicaActa* No.363 pp. 1195–1203(2010)
- [15] A.B. P.Lever, “Inorganic Electronic Spectroscopy” (second edition), Elsevier, Amsterdam p.357,(1984)



- [16] E. T. G. Cavalheiro, Massao Ionashiro, G. Marino, S. T. Breviglieri and G. O. Chierice “Correlation between i.r. spectra and thermal decomposition of cobalt(II), nickel(II), copper(II) and mercury(II) complexes with piperidinedithiocarbamate and pyrrolidinedithiocarbamate” *Transition Metal Chemistry* vol 25, 69-72, (2000).
- [17] Damian C. Onwudiwe and Peter A. Ajibade “Synthesis, Characterization and Thermal Studies of Zn(II), Cd(II) and Hg(II) Complexes of N-Methyl-N-Phenyldithiocarbamate: The Single Crystal Structure of [(C6H5)(CH3)NCS2]4Hg2” *Int. J. Mol. Sci.* 12, 1964-1978,(2011)
- [18] D.Nicholls . “The Chemistry of Iron, Cobalt and Nickel” Pergamon press,Oxford,1st Ed.,1037, 1087, 1088,1090, 1091,1093, 1151,1154, (1973).
- [19] Aysegul Golcu “Transition metal complexes of propranolol dithiocarbamate: synthesis, characterization, analytical properties and biological activity” *Transition Metal Chemistry* vol 31,405–412,(2006)
- [20] A. Manohar, V. Venkatachalam, K. Ramalingam, S. Thirumaran, G. Bocelli, and A. Cantoni “Synthesis, spectral, and single crystal X-ray structural studies on (2,2' bipyridyl) bis(dimethyldithiocarbamato)zinc(II) and (1,10-phenanthroline) bis(dimethyldithio carbamato)zinc(II) ” *Journal of Chemical Crystallography.* Vol. 28, No. 12, (1998)
- [21] R. Pastorek, Z. Travnicek, P.Starha “octahedral nickel(II) hexamethyleneimine-dithiocarbamate complexes involving bidentate N,N-donor ligands” *Inorganica Chimica Acta* 373, 286 -290,(2011)
- [22] K .Nakamoto “Infrared and Raman Spectra of Inorganic and Coordination Compounds”4ed John Wiley & sons ,(1986)